arXiv:0903.3788v1 [math-ph] 23 Mar 2009

A Covering for the dKP–hyper CR Interpolating Equation and Multi-Valued Einstein–Weyl Structures

Oleg I. Morozov

Department of Mathematics, Moscow State Technical University of Civil Aviation,
Kronshtadtskiy Blvd 20, Moscow 125993, Russia
oim@foxcub.org

Abstract. We apply the technique of integrable extensions to the symmetry pseudo-group of the dKP–hyper CR interpolating equation. This allows us to find a covering for this equation and to construct multi-valued Einstein–Weyl structures.

AMS classification scheme numbers: 58H05, 58J70, 35A30

1. Introduction

We consider the equation

$$u_{yy} = u_{tx} + (b\,u_x - c\,u_y)\,u_{xx} + c\,u_x\,u_{xy}, \tag{1}$$

$b = \text{const}$, $c = \text{const}$, found recently by M. Dunajski, [9], as a symmetry reduction of Plebañski's second heavenly equation, [25]. When $c = 0$ and $b \neq 0$, Eq. (1) coincides with the potential form of the Khokhlov–Zabolotskaya equation, [13], also known as dispersionless Kadomtsev–Petviashvili equation (dKP). For $b = 0$ and $c \neq 0$ Eq. (1) becomes the hyperCR equation studied in [24, 8]. The Einstein-Weyl structure associated to (1) is

$$\left\{ \begin{array}{lcl} h & = & (dy - c\,u_x\,dt)^2 - 4\,(dx + (c\,u_y - b\,u_x)\,dt)\,dt, \\ \omega & = & -c\,u_{xx}\,dy + ((c^2\,u_x + 4\,b)\,u_{xx} - 2\,c\,u_{xy})\,dt. \end{array} \right. \tag{2}$$

A Lax pair

$$\left\{ \begin{array}{lcl} q_t & = & ((c\,z - b)\,u_x + c\,u_y + z^2)\; q_x + b\,(z\,u_{xx} + u_{xy})\,q_z, \\ q_y & = & (c\,u_x + z)\,q_x + b\,u_{xx}\,q_z \end{array} \right. \tag{3}$$

is proposed for Eq. (1) in [9]. It contains differentiation with respect to a new independent variable z. The compatibility condition $q_{ty} = q_{yt}$ for system (3) together with the requirement $u_z = 0$ coincides with Eq. (1). The equation $u_z = 0$ does not follow from (3).

When $b \neq 0$ and $c \neq 0$, the simple scaling

$$t = c^4\,b^{-3}\,\tilde{t}, \qquad x = c^2\,b^{-1}\,\tilde{x}, \qquad y = -c^3\,b^{-2}\,\tilde{y} \tag{4}$$

gives after dropping tildes the following equation:

$$u_{yy} = u_{tx} + (u_x + u_y)\,u_{xx} - u_x\,u_{xy}. \tag{5}$$

The aim of this paper is to find a covering, [14, 15, 16, 17], for Eq. (5). We apply the method proposed in [22] and find a contact integrable extension for the Cartan's structure equations of the symmetry pseudo-group of Eq. (5). Integrating the extension yields a Bäcklund transformation and a covering equation for (5). We use these results for constructing multi-valued Einstein–Weyl structures (2) likewise in [21] multi-valued solutions to dKP were found.

2. Preliminaries

2.1. Coverings of PDEs

Let $\pi_\infty : J^\infty(\pi) \to \mathbb{R}^n$ be the infinite jet bundle of local sections of the bundle $\pi : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R}$. The coordinates on $J^\infty(\pi)$ are (x^i, u_I), where $I = (i_1, ..., i_k)$ are symmetric multi-indices, $i_1, ..., i_k \in \{1, ..., n\}$, $u_\emptyset = u$, and for any local section f of π there exists a section $j_\infty(f) : \mathbb{R}^n \to J^\infty(\pi)$ such that $u_I(j_\infty(f)) = \partial^{\#I}(f)/\partial x^{i_1}...\partial x^{i_k}$,

$\#I = \#(i_1, ..., i_k) = k$. The *total derivatives* on $J^\infty(\pi)$ are defined in the local coordintes as

$$D_i = \frac{\partial}{\partial x^i} + \sum_{\#I \geq 0} u_{Ii} \frac{\partial}{\partial u_I}.$$

We have $[D_i, D_j] = 0$ for $i, j \in \{1, ..., n\}$. A differential equation $F(x^i, u_K) = 0$ defines a submanifold $\mathcal{E}^\infty = \{D_I(F) = 0 \mid \#I \geq 0\} \subset J^\infty(\pi)$, where $D_I = D_{i_1} \circ ... \circ D_{i_k}$ for $I = (i_1, ..., i_k)$. We denote restrictions of D_i on $\mathcal{E}^\infty$ as $\bar{D}_i$.

In local coordinates, a *covering* over $\mathcal{E}^\infty$ is a bundle $\widetilde{\mathcal{E}}^\infty = \mathcal{E}^\infty \times \mathcal{V} \to \mathcal{E}^\infty$ with fibre coordinates v^γ, $\gamma \in \{1, ..., N\}$ or $\gamma \in \mathbb{N}$, equipped with *extended total derivatives*

$$\widetilde{D}_i = \bar{D}_i + \sum_\gamma T_i^\gamma(x^j, u_I, v^\beta) \frac{\partial}{\partial v^\gamma}$$

such that $[\widetilde{D}_i, \widetilde{D}_j] = 0$ whenever $(x^i, u_I) \in \mathcal{E}^\infty$. Action of $\widetilde{D}_i$ on the fibre variables v^γ gives

$$v^\gamma_{x^i} = T_i^\gamma(x^j, u_I, v^\beta).$$

These equations are called a *Bäcklund transformation*. Excluding u from them yields a system of PDEs for v^γ. This system is called a *covering equation*.

In terms of differential forms, the covering is defined by the forms, [28],

$$\omega^\gamma = dv^\gamma - T_i^\gamma(x^j, u_I, v^\beta)\, dx^i$$

such that

$$d\omega^\gamma \equiv 0 \mod \omega^\beta, \bar{\vartheta}_I \quad \Longleftrightarrow \quad (x^i, u_I) \in \mathcal{E}^\infty, \tag{6}$$

where $\bar{\vartheta}_I$ are restrictions of contact forms $\vartheta_I = du_I - u_{I,k}\, dx^k$ on $\mathcal{E}^\infty$.

2.2. Cartan's structure theory of Lie pseudo-groups

Let M be a manifold of dimension n. A *local diffeomorphism* on M is a diffeomorphism $\Phi : \mathcal{U} \to \hat{\mathcal{U}}$ of two open subsets of M. A *pseudo-group* $\mathfrak{G}$ on M is a collection of local diffeomorphisms of M, which is closed under composition *when defined*, contains an identity and is closed under inverse. A *Lie pseudo-group* is a pseudo-group whose diffeomorphisms are local analytic solutions of an involutive system of partial differential equations.

Élie Cartan's approach to Lie pseudo-groups is based on a possibility to characterize transformations from a pseudo-group in terms of a set of invariant differential 1-forms called *Maurer–Cartan* (MC) *forms*. In a general case MC forms ω^1, ..., ω^m of a Lie pseudo-group $\mathfrak{G}$ on M are defined on a direct product $\tilde{M} \times G$, where $\mu : \tilde{M} \times G \to M$ is a bundle, $m = \dim \tilde{M}$, G is a finite-dimensional Lie group. The forms ω^i are semi-basic w.r.t. the natural projection $\tilde{M} \times G \to \tilde{M}$ and define a coframe on $\tilde{M}$, that is, a basis of the cotangent bundle of $\tilde{M}$. They characterize the pseudo-group $\mathfrak{G}$ in the following sense: a local diffeomorphism $\Phi : \mathcal{U} \to \hat{\mathcal{U}}$ on M belongs to $\mathfrak{G}$ whenever there exists a local diffeomorphism $\Psi : \mathcal{V} \to \hat{\mathcal{V}}$ on $\tilde{M} \times G$ such that $\mu \circ \Psi = \Phi \circ \mu$ and the forms ω^j are invariant w.r.t. Ψ, that is,

$$\Psi^* \left(\omega^i|_{\hat{\mathcal{V}}}\right) = \omega^i|_{\mathcal{V}}. \tag{7}$$

Expressions of exterior differentials of the forms ω^i in terms of themselves give Cartan's structure equations of $\mathfrak{G}$:

$$d\omega^i = A^i_{\gamma j}\, \pi^\gamma \wedge \omega^j + B^i_{jk}\, \omega^j \wedge \omega^k, \qquad B^i_{jk} = -B^i_{kj}. \tag{8}$$

Here and below we assume summation on repeated indices. The forms π^γ, $\gamma \in \{1, ..., \dim G\}$, are linear combinations of MC forms of the Lie group G and the forms ω^i. The coefficients $A^i_{\gamma j}$ and B^i_{jk} are either constants or functions of a set of invariants $U^\kappa : M \to \mathbb{R}$, $\kappa \in \{1, ..., l\}$, $l < \dim M$, of the pseudo-group $\mathfrak{G}$, so $\Phi^* \left(U^\kappa|_{\hat{\mathcal{U}}}\right) = U^\kappa|_{\mathcal{U}}$ for every $\Phi \in \mathfrak{G}$. In the latter case, differentials of U^κ are invariant 1-forms, so they are linear combinations of the forms ω^j,

$$dU^\kappa = C^\kappa_j\, \omega^j, \tag{9}$$

where the coefficients C^κ_j depend on the invariants U^1, ..., U^l only.

Eqs. (8) must be compatible in the following sense: we have

$$d(d\omega^i) = 0 = d\left(A^i_{\gamma j}\, \pi^\gamma \wedge \omega^j + B^i_{jk}\, \omega^j \wedge \omega^k\right), \tag{10}$$

therefore there must exist expressions

$$d\pi^\gamma = W^\gamma_{\lambda j}\, \chi^\lambda \wedge \omega^j + X^\gamma_{\beta\epsilon}\, \pi^\beta \wedge \pi^\epsilon + Y^\gamma_{\beta j}\, \pi^\beta \wedge \omega^j + Z^\gamma_{jk}\, \omega^j \wedge \omega^k, \tag{11}$$

with some additional 1-forms χ^λ and the coefficients $W^\gamma_{\lambda j}$ to Z^γ_{jk} depending on the invariants U^κ such that the right-hand side of (10) appear to be identically equal to zero after substituting for (8), (9), and (11). Also, from (9) it follows that the right-hand side of the equation

$$d(dU^\kappa) = 0 = d(C^\kappa_j\, \omega^j) \tag{12}$$

must be identically equal to zero after substituting for (8) and (9).

The forms π^γ are not invariant w.r.t. the pseudo-group $\mathfrak{G}$. Respectively, the structure equations (8) are not changing when replacing $\pi^\gamma \mapsto \pi^\gamma + z^\gamma_j\, \omega^j$ for certain parametric coefficients z^γ_j. The dimension $r^{(1)}$ of the linear space of these coefficients satisfies the following inequality

$$r^{(1)} \le n \dim G - \sum_{k=1}^{n-1} (n-k)\, \sigma_k, \tag{13}$$

where the *reduced characters* σ_k are defined by

$$\sigma_k = \max_{u_1, ..., u_k} \operatorname{rank} \mathbb{A}_k(u_1, ..., u_k)$$

with the matrices $\mathbb{A}_k$ inductively defined by

$$\mathbb{A}_1(u_1) = \left(A^i_{\gamma j}\, u^j_1\right), \qquad \mathbb{A}_l(u_1, ..., u_l) = \left(\begin{array}{c} \mathbb{A}_{l-1}(u_1, ..., u_{l-1}) \\ A^i_{\gamma j}\, u^j_l \end{array}\right),$$

see [4, §5], [23, Def. 11.4] for the full discussion. The system of forms ω^k is *involutive* if (13) is an equality, [4, §6], [23, Def. 11.7].

Cartan's fundamental theorems, [4, §§16, 22–24], [7], [27, §§16, 19, 20, 25,26], [26, §§14.1–14.3], state that for a Lie pseudo-group there exists a set of MC forms whose structure equations satisfy the compatibility and involutivity conditions; conversely, if Eqs. (8), (9) meet the compatibility conditions (10), (12), and the involutivity condition, then there exists a collection of 1-forms $\omega^1, \dots, \omega^m$ and functions $U^1, \dots, U^l$ which satisfy (8) and (9). Eqs. (7) then define local diffeomorphisms from a Lie pseudo-group.

Example 1. Consider the bundle $J^2(\pi)$ of jets of the second order of the bundle $\pi : \mathbb{R}^n \times \mathbb{R} \to \mathbb{R}^n$, $\pi : (x^1, ..., x^n, u) \mapsto (x^1, ..., x^n, u)$. A differential 1-form ϑ on $J^2(\pi)$ is called a *contact form* if it is annihilated by all 2-jets of local sections f of the bundle π: $j_2(f)^*\vartheta = 0$. In the local coordinates every contact 1-form is a linear combination of the forms $\vartheta_0 = du - u_i\,dx^i$, $\vartheta_i = du_i - u_{ij}\,dx^j$, $i, j \in \{1, ..., n\}$, $u_{ji} = u_{ij}$. A local diffeomorphism $\Delta : J^2(\pi) \to J^2(\pi)$, $\Delta : (x^i, u, u_i, u_{ij}) \mapsto (\hat{x}^i, \hat{u}, \hat{u}_i, \hat{u}_{ij})$, is called a *contact transformation* if for every contact 1-form $\hat{\vartheta}$ the form $\Delta^*\hat{\vartheta}$ is also contact. We denote by $\mathrm{Cont}(J^2(\pi))$ the pseudo-group of contact transformations on $J^2(\pi)$. Consider the following 1-forms

$$\begin{aligned}
&\Theta_0 = a\,\vartheta_0, \quad \Theta_i = g_i\,\Theta_0 + a\,B^k_i\,\vartheta_k, \quad \Xi^i = c^i\,\Theta_0 + f^{ik}\,\Theta_k + b^i_k\,dx^k,\\
&\Theta_{ij} = a\,B^k_i\,B^l_j\,(du_{kl} - u_{klm}\,dx^m) + s_{ij}\,\Theta_0 + w^k_{ij}\,\Theta_k,
\end{aligned} \tag{14}$$

defined on $J^2(\pi) \times \mathcal{H}$, where $\mathcal{H}$ is an open subset of $\mathbb{R}^{(2n+1)(n+3)(n+1)/3}$ with local coordinates $(a,\ b^i_k,\ c^i,\ f^{ik},\ g_i,\ s_{ij},\ w^k_{ij},\ u_{ijk})$, $i, j, k, \in \{1, ..., n\}$, such that $a \neq 0$, $\det(b^i_k) \neq 0$, $f^{ik} = f^{ki}$, $s_{ij} = s_{ji}$, $w^k_{ij} = w^k_{ji}$, $u_{ijk} = u_{ikj} = u_{jik}$, while (B^i_k) is the inverse matrix for the matrix (b^i_k). As it is shown in [20], the forms (14) are MC forms for $\mathrm{Cont}(J^2(\pi))$, that is, a local diffeomorphism $\widetilde{\Delta} : J^2(\pi) \times \mathcal{H} \to J^2(\pi) \times \mathcal{H}$ satisfies the conditions $\widetilde{\Delta}^*\,\hat{\Theta}_0 = \Theta_0$, $\widetilde{\Delta}^*\,\hat{\Theta}_i = \Theta_i$, $\widetilde{\Delta}^*\,\hat{\Xi}^i = \Xi^i$, and $\widetilde{\Delta}^*\,\hat{\Theta}_{ij} = \Theta_{ij}$ if and only if it is projectable on $J^2(\pi)$, and its projection $\Delta : J^2(\pi) \to J^2(\pi)$ is a contact transformation. The structure equations for $\mathrm{Cont}(J^2(\pi))$ have the form

$$\begin{aligned}
d\Theta_0 &= \Phi^0_0 \wedge \Theta_0 + \Xi^i \wedge \Theta_i,\\
d\Theta_i &= \Phi^0_i \wedge \Theta_0 + \Phi^k_i \wedge \Theta_k,\\
d\Xi^i &= \Phi^0_0 \wedge \Xi^i - \Phi^i_k \wedge \Xi^k + \Psi^{i0} \wedge \Theta_0 + \Psi^{ik} \wedge \Theta_k,\\
d\Theta_{ij} &= \Phi^k_i \wedge \Theta_{kj} + \Phi^k_j \wedge \Theta_{ki} - \Phi^0_0 \wedge \Theta_{ij} + \Upsilon^0_{ij} \wedge \Theta_0 + \Upsilon^k_{ij} \wedge \Theta_k + \Xi^k \wedge \Theta_{ijk},
\end{aligned}$$

where the additional forms Φ^0_0, Φ^0_i, Φ^k_i, Ψ^{i0}, Ψ^{ij}, Υ^0_{ij}, Υ^k_{ij}, and Θ_{ijk} depend on differentials of the coordinates of $\mathcal{H}$.

Example 2. Suppose $\mathcal{E}$ is a second-order differential equation in one dependent and n independent variables. We consider $\mathcal{E}$ as a submanifold in $J^2(\pi)$. Let $\mathrm{Cont}(\mathcal{E})$ be the group of contact symmetries for $\mathcal{E}$. It consists of all the contact transformations on $J^2(\pi)$ mapping $\mathcal{E}$ to itself. Let $\iota_0 : \mathcal{E} \to J^2(\pi)$ be an embedding, and $\iota = \iota_0 \times \mathrm{id} : \mathcal{E} \times \mathcal{H} \to \mathrm{J}^2(\pi) \times \mathcal{H}$. The MC forms of $\mathrm{Cont}(\mathcal{E})$ can be computed from the forms $\theta_0 = \iota^*\Theta_0$, $\theta_i = \iota^*\Theta_i$, $\xi^i = \iota^*\Xi^i$, and $\theta_{ij} = \iota^*\Theta_{ij}$ algorithmically by means of Cartan's method of equivalence, [4, 5, 6, 7, 10, 12, 23], see details and examples in [11, 19, 20].

3. Cartan's structure of the symmetry pseudo-group for the dKP-hyperCR interpolating equation

We use the method outlined in the previous section to compute MC forms and structure equations of the pseudo-group of contact symmetries for Eq. (5). We write out here the structure equations for the forms θ_0, θ_j, ξ^j with $j \in \{1,2,3\}$ only; the full set of the structure equations is given in Appendix.

We have

$$\begin{aligned}
d\theta_0 &= \eta_1 \wedge \theta_0 + \xi_1 \wedge \theta_1 + \xi_2 \wedge \theta_2 + \xi_1 \wedge \theta_1 + \xi_3 \wedge \theta_3,\\
d\theta_1 &= \tfrac{1}{2}\,\eta_1 \wedge \theta_1 + \tfrac{1}{4}\,\eta_2 \wedge (\theta_0 + 8\,\theta_3) + \tfrac{1}{16}\,\left(24\,(\theta_{22} - U\,\xi_1 - \xi_2) - 5\,\xi_3\right) \wedge \theta_1 + 2\,\theta_2 \wedge \theta_3\\
&\quad + \tfrac{1}{8}\,\left(11\,\theta_2 - 16\,V\,\theta_{22} + 8\,\theta_{23} + 4\,(2\,V - 1)\,\xi_2 + (8\,V - 3\,U)\,\xi_3\right) \wedge \theta_0 + \xi_1 \wedge \theta_{11}\\
&\quad + \xi_2 \wedge \theta_{12} + \xi_3 \wedge \theta_{13},\\
d\theta_2 &= \tfrac{1}{16}\,\left(8\,(\eta_1 + \theta_{22} - U\,\xi_1 - \xi_2) - 3\,\xi_3\right) \wedge \theta_2 + \xi_1 \wedge \theta_{12} + \xi_2 \wedge \theta_{22} + \xi_3 \wedge \theta_{23},\\
d\theta_3 &= \eta_2 \wedge \theta_2 + \left(\eta_1 + \theta_{22} - U\,\xi_1 - \xi_2 - \tfrac{1}{4}\,\xi_3\right) \wedge \theta_3 + \tfrac{5}{64}\,(8\,\theta_{22} - \xi_2 + 3\,\xi_3) \wedge \theta_0 + \xi_1 \wedge \theta_{13}\\
&\quad + \xi_2 \wedge \theta_{23} + \xi_3 \wedge \theta_{12},\\
d\xi_1 &= \; - \tfrac{1}{16}\,\left(8\,\eta_1 + 24\,(\,\theta_{22} + \xi_2) - 5\,\xi_3\right) \wedge \xi_1,\\
d\xi_2 &= \tfrac{1}{8}\,(5\,\theta_0 + 8\,V\,\theta_2 + 8\,\theta_3 - 4\,U\,\xi_2) \wedge \xi_1 + \tfrac{1}{16}\,\left(8\,(\eta_1 - \theta_{22}) + 3\,\xi_3\right) \wedge \xi_2 - \eta_2 \wedge \xi_3,\\
d\xi_3 &= \; - (2\,\eta_2 + \theta_2 + U\,\xi_3) \wedge \xi_1 - (\theta_{22} - \xi_2) \wedge \xi_3,
\end{aligned}$$

where the invariants

$$\begin{aligned}
U &= u_{xx}^{-4}\,\left((u_x + u_y)\,u_{xxx}^2 + (u_{txx} - u_x\,u_{xxy} + u_{xx}\,(u_{xy} + 4\,u_{xx})\,u_{xxx} - u_{xxy}^2 - u_{xx}^2\,u_{xxy}\right),\\
V &= u_{xxx}\,u_{xx}^{-2}
\end{aligned}$$

satisfy

$$\begin{aligned}
dU &= \tfrac{1}{2}\,U\,\eta_1 + \eta_2 + \eta_3 - \tfrac{5}{8}\,\theta_0 - \left(V + \tfrac{1}{8}\right)\,\theta_2 - \theta_3 - \left(4\,V - \tfrac{3}{2}\,U\right)\,\theta_{22} - \theta_{23}\\
&\quad + \left(4\,V - U + \tfrac{1}{2}\right)\,\xi_2 - \left(V - \tfrac{11}{16}\,U\right)\,\xi_3, && (15)\\
dV &= \tfrac{1}{2}\,V\,\left(\eta_1 + \theta_{22} + (6\,V - U)\,\xi_1 - \xi_2\right) - \tfrac{5}{16}\,V\,\xi_3, && (16)
\end{aligned}$$

and where

$$\begin{aligned}
\theta_0 &= V^2\,u_{xx}^3\,(du - u_t\,dt - u_x\,dx - u_y\,dy),\\
\theta_2 &= \; -V\,\left(du_x - u_{tx}\,dt - u_{xx}\,dx - u_{xy}\,dy\right),\\
\theta_3 &= V\,u_{xxy}\,u_{xx}^{-2}\,du_x - V^2\,du_y + V\,(V\,u_{ty} - u_{tx}\,u_{xxy}\,u_{xx}^{-2})\,dt + V\,(V\,u_{xy} - u_{xxy}\,u_{xx}^{-1})\,dx\\
&\quad + V\,\left(V\,(u_{tx} + (u_x + u_y)\,u_{xx} - u_x\,u_{xy}) - u_{xy}\,u_{xxy}\,u_{xx}^{-2}\right)\,dy - \tfrac{5}{8}\,\theta_0,\\
\theta_{22} &= \; -u_{xx}^{-1}\,\left(du_{xx} - u_{txx}\,dt - u_{xxx}\,dx - u_{xxy}\,dy\right),\\
\xi_1 &= u_{xx}\,V^{-1}\,dt, && (17)\\
\xi_2 &= u_{xx}^{-1}\,\left(u_{xxx}\,dx + u_{xxy}\,dy - (u_{xxx}\,(u_x + u_y) - u_{xxy}\,u_x - u_{xxy}^2\,u_{xxx}^{-1})\,dt\right),\\
\xi_3 &= u_{xx}\,(u_x + 2\,u_{xxy}\,u_{xxx}^{-1})\,dt + u_{xx}\,dy,\\
\eta_1 &= 2\,u_{xxx}^{-1}\,du_{xxx} + \left(10\,V - 3\,U + 4\,(V\,u_{xy}\,u_{xx}^{-1} - u_{xxy}\,u_{xx}^{-2})\right)\,\xi_1 + \tfrac{5}{8}\,\xi_3 + 3\,(\theta_{22} - \xi_2),\\
\eta_2 &= \; -u_{xx}^{-2}\,du_{xxy} - \left(V^2\,(u_x + u_y + (u_{tx} - u_{xy}\,(u_x - 1))\,u_{xx}^{-2}\right)
\end{aligned}$$

$$
\begin{aligned}
&+ \big(4\,V\,u_{xxy} - \tfrac{3}{2}\,U\,u_{xxy}\big)\;u_{xx}^{-2} - u_{xxy}^{2}\,u_{xx}^{-4}\big)\;\xi_1 + \tfrac{1}{2}\,(3\,u_{xxy}\,u_{xx}^{-2} - 1)\,\xi_2 \\
&- \big(V\,(u_{xy}\,u_{xx} + 2) - \tfrac{11}{16}\,u_{xxy}\,u_{xx}^{-2}\big)\;\xi_3 - \tfrac{3}{2}\,u_{xxy}\,u_{xx}^{-2}\,\theta_{22} + \tfrac{1}{2}\,u_{xxy}\,u_{xx}^{-2}\,\eta_1.
\end{aligned}
$$

4. Integrable extensions

In [3, §6], the definition of integrable extension of an exterior differential system is designed to study finite-dimensional coverings. In general case, coverings of PDEs with three or more independent variables are infinite-dimensional, [18]. To cope with infinite-dimensional coverings we use a natural generalization of the definition, [22].

Suppose $\mathfrak{G}$ is a Lie pseudo-group on a manifold M and $\omega^1, \dots, \omega^m$ are its MC forms with structure equations (8), (9). Consider a system of equations

$$
\begin{aligned}
d\tau^q &= D^q_{\rho r}\,\eta^\rho \wedge \tau^r + E^q_{rs}\,\tau^r \wedge \tau^s + F^q_{r\beta}\,\tau^r \wedge \pi^\beta + G^q_{rj}\,\tau^r \wedge \omega^j + H^q_{\beta j}\,\pi^\beta \wedge \omega^j \\
&\quad + I^q_{jk}\,\omega^j \wedge \omega^k, \qquad (18)\\
dV^\epsilon &= J^\epsilon_j\,\omega^j + K^\epsilon_q\,\tau^q, \qquad (19)
\end{aligned}
$$

with unknown 1-forms τ^q, $q \in \{1, \dots, Q\}$, η^ρ, $\rho \in \{1, \dots, R\}$, and unknown functions V^ϵ, $\epsilon \in \{1, \dots, S\}$ for some $Q, R, S \in \mathbb{N}$. The coefficients $D^q_{\rho r}$, ..., K^ϵ_q in (18), (19) are supposed to be functions of U^q and V^γ.

Definition 1. System (18), (19) is an *integrable extension* of system (8), (9), if Eqs. (18), (19), (8), and (9) together satisfy the compatibility and involutivity conditions.

In this case from Cartan's third fundamental theorem for Lie pseudo-groups it follows that there exists a set of forms τ^q and functions V^ϵ which are solutions to Eqs. (18) and (19). Then τ^q, V^ϵ together with ω^i, U^q define a Lie pseudo-group on a manifold $N \cong M \times \mathbb{R}^Q$.

Definition 2. The integrable extension is called *trivial*, if there is a change of variables on N such that in the new variables the coefficients $F^q_{r\beta}$, G^q_{rj}, $H^q_{\beta j}$, I^q_{jk}, and J^ϵ_j are equal to zero, while the coefficients $D^q_{\rho r}$, E^q_{rs}, and K^ϵ_q are independent of U^q. Otherwise, the integrable extension is called *non-trivial.*

Let θ^γ_I and ξ^j be a set of MC forms of the symmetry pseudo-group $\mathrm{Cont}(\mathcal{E})$ of a PDE $\mathcal{E}$ such that $\xi^1 \wedge \dots \wedge \xi^n \neq 0$ on any solution manifold of $\mathcal{E}$, while θ^γ_I are contact forms. We take the following reformulation of the definition (6) of a covering.

Definition 3. A non-trivial integrable extension of the form

$$
d\omega^q = \Pi^q_r \wedge \omega^r + \xi^j \wedge \Omega^q_j \qquad (20)
$$

is called a *contact integrable extension* (CIE) of the structure equations of $\mathrm{Cont}(\mathcal{E})$ if

1. Π^q_r are some non-trivial differential 1-forms,
2. $\Omega^q_j \equiv 0 \mod \theta^\gamma_I, \omega^q_j$ for some additional 1-forms ω^q_j,
3. $\Omega^q_j \not\equiv 0 \mod \omega^q_j$.

Since (20) is integrable extension, Cartan's theorem yields existence of the forms ω^q satisfying (20). From [2, Ch. IV, Prop. 5.10] it follows that the forms ω^q define a system of PDEs. This system is a covering for $\mathcal{E}$.

We apply this construction to the structure equations (35), (15), and (16) of the symmetry pseudo-group of Eq. (5). We restrict our analysis to CIEs of the form

$$
\begin{aligned}
d\omega_0 = {} & \left(\sum_{i=0}^{3} A_i \,\theta_i + \sum{}^{*} B_{ij}\,\theta_{ij} + \sum_{s=1}^{7} C_s\,\eta_s + \sum_{j=1}^{3} D_j\,\xi^j + E\,\omega_1 \right) \wedge \omega_0 \\
& + \sum_{j=1}^{3} \left(\sum_{k=0}^{3} F_{jk}\,\theta_k + G_j\,\omega_1 \right) \wedge \xi^j, \qquad (21)
\end{aligned}
$$

where $\sum{}^{*}$ means summation for all $i, j \in \mathbb{N}$ such that $1 \le i \le j \le 3$, $(i,j) \neq (3,3)$, and consider two types of these CIEs:

TYPE 1 — the coefficients A_i to G_j in (21) depend on the invariants U, V of the symmetry pseudo-group of Eq. (5) only;

TYPE 2 — the coefficients A_i to G_j depend also on one additional invariant, say W. In this case, the differential of this new invariant satisfies the following equation

$$
dW = \sum_{i=0}^{3} H_i\,\theta_i + \sum{}^{*} I_{ij}\,\theta_{ij} + \sum_{s=1}^{7} J_s\,\eta_s + \sum_{j=1}^{3} K_j\,\xi^j + \sum_{q=0}^{1} L_q\,\omega_q, \qquad (22)
$$

where the coefficients H_i to L_q are functions of U, V, W.

The requirements of Defintions 1 and 3 yield over-determined systems for the coefficients A_i to G_j of the CIE of the first type and A_i to L_q of the CIE of the second type. The results of analysis of these systems are summarized in the following theorem.

Theorem 1. *There is no a* CIE *of the first type for the structure equations (35), (15), and (16). Every their* CIE *of the second type is contact-equivalent to the following one:*

$$
\begin{aligned}
d\omega_0 = {} & \left(\tfrac{1}{2}\,(\eta_1 - \theta_{22}) - \omega_1 - \tfrac{1}{2}(W^2 + 2\,(V - W) - U)\,\xi_1 - \tfrac{1}{16}\,(8\,W - 11)\,\xi_3 \right) \wedge \omega_0 \\
& + \left(W^2\,\omega_1 + \tfrac{5}{8}\,\theta_0 + W\,\theta_2 + \theta_3 \right) \wedge \xi_1 + \omega_1 \wedge \xi_2 + (W\,\omega_1 + \theta_2) \wedge \xi_3, \qquad (23)
\end{aligned}
$$

$$
\begin{aligned}
dW = {} & V\,\omega_1 + \tfrac{1}{2}\,W\,\eta_1 + \eta_2 + \tfrac{1}{2}\,W\,\theta_{22} + \theta_2 + \tfrac{1}{2}\,W\,(V\,(W+4) - U)\,\xi_1 \\
& + \tfrac{1}{2}\,(V - W - 1)\,\xi_2 + \tfrac{1}{16}\,(W\,(4\,V - 5) + 8\,V)\,\xi_3. \qquad (24)
\end{aligned}
$$

Since the forms (17) are known, it is easy to find the form ω_0 explicitly:

Theorem 2. *We have the following solution to Eq. (23) up to a contact equivalence:*

$$
\begin{aligned}
\omega_0 = {} & u_{xxx}\,u_{xx}^{-1}\,v_x^{-1} \left(dv - v_x \left(\ln^2|v_x| - u_y - (\ln|v_x| + 1)\,u_x + 1 \right) dt \right. \\
& \left. - v_x\,dx - v_x\,(\ln|v_x| - u_x)\,dy \right). \qquad (25)
\end{aligned}
$$

This form defines the following Bäcklund transformation or a Lax pair:

$$
\left\{
\begin{array}{lcl}
v_t & = & v_x \left(\ln^2|v_x| - u_y - (\ln|v_x| + 1)\,u_x + 1 \right), \\
v_y & = & v_x\,(\ln|v_x| - u_x).
\end{array}
\right. \qquad (26)
$$

It is easy to verify directly that Eqs. (26) are compatible whenever Eq. (5) is satisfied. From Eqs. (26) it follows that

$$
\left\{
\begin{array}{lcl}
u_x & = & \ln|v_x| - v_y\,v_x^{-1}, \\
u_y & = & (v_y\,(\ln|v_x| + 1) - v_t)\;v_x^{-1} - \ln|v_x| + 1,
\end{array}
\right. \qquad (27)
$$

Cross-differentiating u in this system yields the covering equation:

$$v_{yy} = v_{tx} + \big((v_y \, \ln|v_x| - v_t)\, v_x^{-1} + 1\big) \; v_{xx} + \big(v_y \, v_x^{-1} - \ln|v_x|\big) \; v_{xy}. \tag{28}$$

5. Multi-valued Einstein–Weyl Structures

We use the results of the previous section to construct a family of Einstein–Weyl structures (2) depending on two arbitrary functions of one variable. We take the ansatz, [1, Ch. VIII, § 5.IV], [21],

$$v_t = F(v_x), \qquad v_y = G(v_x). \tag{29}$$

This system is compatible for every (smooth) functions F and G. Substituting for (29) into (28) and denoting

$$v_x = s \tag{30}$$

yields

$$(G'(s))^2 = F'(s) + (G(s) \, \ln|s| - F(s)) \; s^{-1} + 1 + \big(G(s)\, s^{-1} - \ln|s|\big) \; G'(s).$$

We consider this as an ODE for the unknown fucntion F and the functional parameter G being an arbitrary smooth function. Then we have

$$F(s) = s \int s^{-1} \; \Big((G'(s))^2 + \ln|s| \, (G'(s) - s^{-1}\, G(s)) - s^{-1}\, G(s)\, G'(s) - 1\Big) \; ds. \tag{31}$$

From (29) and (30) it follows that the function s satisfies the compatible system of PDEs

$$s_t = F'(s)\, s_x, \qquad s_y = G'(s)\, s_x. \tag{32}$$

The general solution of this system in the implicit form reads

$$s = Q(x + t\, F'(s) + y\, G'(s)), \tag{33}$$

where Q is an arbitrary (smooth) function of one variable. For $t = 0$ and $y = 0$ we have $s = Q(x)$, so Q is an initial value for Eqs. (32). In general, Eq. (33) defines s as a multi-valued function of t, x, and y.

Then Eqs. (27), (29), and (30) give

$$\left\{ \begin{array}{lcl} u_x & = & \ln|s| - s^{-1}\, G(s), \\ u_y & = & (s^{-1}\, G(s) - 1) \, \ln|s| - s^{-1}\, F(s) + 1, \end{array} \right. \tag{34}$$

where the function F is defined by Eq. (31), and the function s is defined by (33). Eqs. (34) together with the scaling (4) provide a family of Einstein-Weyl structures (2) depending on two arbitrary functions of one variable.

Figures 1 to 4 show graphs of u_x and u_y at $t = -10$ and $t = 10$ for the choice of $G(s) = -3\, s$, $Q(x) = x^2 + 1$, and $F(s) = (5 - \ln|s|)\, s$.



Figure 1. The graph of u_x at $t = -10$.



Figure 2. The graph of u_y at $t = -10$.



Figure 3. The graph of u_x at $t = 10$.



Figure 4. The graph of u_y at $t = 10$.

Acknowledgements

I am grateful to the National Defense University of Taiwan, where the part of this work was done. My special thanks are to Jen-Hsu Chang for the warm hospitality in the NDU and valuable discussions, and to M.V. Pavlov for organizing my visit to Taiwan. The work was partially supported by the Russian-Taiwanese grant 95WFE0300007 (RFBR grant 06-01-89507-HHC) and the joint grant 09-01-92438-KE_a of RFBR (Russia) and Consortium E.I.N.S.T.E.IN (Italy).

Appendix

The structure equations of the symmetry pseudo-group for the dKP-hyperCR interpolating equation read

$$
\begin{aligned}
d\theta_0 &= \eta_1 \wedge \theta_0 + \xi_1 \wedge \theta_1 + \xi_2 \wedge \theta_2 + \xi_1 \wedge \theta_1 + \xi_3 \wedge \theta_3, \\
d\theta_1 &= \tfrac{1}{2}\,\eta_1 \wedge \theta_1 + \tfrac{1}{4}\,\eta_2 \wedge (\theta_0 + 8\,\theta_3) + \tfrac{1}{16}\,(24\,(\theta_{22} - U\,\xi_1 - \xi_2) - 5\,\xi_3) \wedge \theta_1 + 2\,\theta_2 \wedge \theta_3 \\
&\quad + \tfrac{1}{8}\,(11\,\theta_2 - 16\,V\,\theta_{22} + 8\,\theta_{23} + 4\,(2\,V - 1)\,\xi_2 + (8\,V - 3\,U)\,\xi_3) \wedge \theta_0 + \xi_1 \wedge \theta_{11} \\
&\quad + \xi_2 \wedge \theta_{12} + \xi_3 \wedge \theta_{13}, \\
d\theta_2 &= \tfrac{1}{16}\,(8\,(\eta_1 + \theta_{22} - U\,\xi_1 - \xi_2) - 3\,\xi_3) \wedge \theta_2 + \xi_1 \wedge \theta_{12} + \xi_2 \wedge \theta_{22} + \xi_3 \wedge \theta_{23}, \\
d\theta_3 &= \eta_2 \wedge \theta_2 + \left(\eta_1 + \theta_{22} - U\,\xi_1 - \xi_2 - \tfrac{1}{4}\,\xi_3\right) \wedge \theta_3 + \tfrac{5}{64}\,(8\,\theta_{22} - \xi_2 + 3\,\xi_3) \wedge \theta_0 + \xi_1 \wedge \theta_{13} \\
&\quad + \xi_2 \wedge \theta_{23} + \xi_3 \wedge \theta_{12}, \\
d\xi_1 &= -\,\tfrac{1}{16}\,(8\,\eta_1 + 24\,(\,\theta_{22} + \xi_2) - 5\,\xi_3) \wedge \xi_1, \\
d\xi_2 &= \tfrac{1}{8}\,(5\,\theta_0 + 8\,V\,\theta_2 + 8\,\theta_3 - 4\,U\,\xi_2) \wedge \xi_1 + \tfrac{1}{16}\,(8\,(\eta_1 - \theta_{22}) + 3\,\xi_3) \wedge \xi_2 - \eta_2 \wedge \xi_3, \\
d\xi_3 &= -\,(2\,\eta_2 + \theta_2 + U\,\xi_3) \wedge \xi_1 - (\theta_{22} - \xi_2) \wedge \xi_3, \\
d\theta_{11} &= \tfrac{3}{2}\,\eta_2 \wedge \theta_1 + \left(\left(2\,V - \tfrac{1}{2}\,U\right)\,\eta_2 + 2\,V\,\eta_3 - \eta_4\right) \wedge \theta_0 + \eta_5 \wedge \xi_2 + \eta_6 \wedge \xi_3 + \eta_7 \wedge \xi_1 \\
&\quad + \left(\left(U - 2\,V^2 + \tfrac{51}{32}\,V\right)\theta_2 + (1 - 2\,V)\,\theta_3 + \tfrac{7}{4}\,\theta_{12} - 2\,V\,(U + 3\,V)\,\theta_{22}\right. \\
&\quad + \tfrac{1}{4}\,(6\,U - V)\,\theta_{23} + \tfrac{1}{8}\,(U\,(24\,V - 7) + V\,(48\,V - 1))\,\xi_2 \\
&\quad \left. + \tfrac{1}{16}\,\left(44\,V^2 - 15\,U^2 + 28\,U\,V\right)\xi_3\right) \wedge \theta_0 + \left(2\,\theta_2 - 11\,V\,\theta_{22} + \theta_{23}\right. \\
&\quad \left. + \tfrac{1}{2}\,(22\,V - 1)\,\xi_2 + \tfrac{1}{4}\,(4\,V + U)\,\xi_3\right) \wedge \theta_1 + (V\,\theta_{12} - 3\,\theta_{13}) \wedge \theta_2 \\
&\quad + \left(\tfrac{3}{4}\,V\,\theta_2 + \theta_{12} - 2\,V\,\theta_{23} + (U - 2\,V)\,\xi_2 - 2\,V\,(U + V)\,\xi_3\right) \wedge \theta_3 \\
&\quad + \left(2\,\eta_1 + 3\,\theta_{22} - 3\,\xi_2 - \tfrac{5}{8}\,\xi_3\right) \wedge \theta_{11} + 2\,U\,\xi_2 \wedge \theta_{12} + \left(4\,\eta_2 + \tfrac{5}{2}\,U\,\xi_3\right) \wedge \theta_{13}, \\
d\theta_{12} &= \left(\tfrac{1}{4}\,\eta_2 - 2\,V\,\theta_{22} - \theta_{23} + 3\,V\,\xi_2 + \tfrac{1}{8}\,U\,\xi_3\right) \wedge \theta_2 + \eta_3 \wedge \xi_2 + \eta_4 \wedge \xi_3 + \eta_5 \wedge \xi_1 \\
&\quad + \left(\eta_1 + 2\,\theta_{22} - 2\,\xi_2 - \tfrac{1}{2}\,\xi_3\right) \wedge \theta_{12} - \left(\tfrac{5}{8}\,\theta_0 - \theta_3 + U\,\xi_2\right) \wedge \theta_{22} \\
&\quad + \left(2\,\eta_2 + \tfrac{3}{2}\,U\,\xi_3\right) \wedge \theta_{23}, \\
d\theta_{13} &= \tfrac{5}{64}\,(6\,\eta_2 - 8\,(\eta_3 - \theta_3 - U\,\theta_{22} - \theta_{23}) + (8\,V - 7)\,\theta_2 - 6\,U\,(2\,\xi_2 + \xi_3)) \wedge \theta_0 + \eta_4 \wedge \xi_2 \\
&\quad + \eta_5 \wedge \xi_3 + \eta_6 \wedge \xi_1 + \tfrac{5}{64}\,(8\,(\theta_{22} - \xi_2) + 3\,\xi_3) \wedge \theta_1 + (3\,\eta_2 + 2\,\theta_2 + 2\,U\,\xi_3) \wedge \theta_{12} \\
&\quad - \left(\tfrac{7}{8}\,\theta_3 + \tfrac{1}{2}\,(2\,V - U)\,\xi_2 + V\,(U + V)\,\xi_3\right) \wedge \theta_2 - \left(\theta_3 - \tfrac{3}{2}\,U\,\xi_2\right) \wedge \theta_{23} \\
&\quad + \tfrac{3}{8}\,(2\,\eta_2 - 4\,V\,(\theta_{22} - \xi_2) + U\,\xi_3) \wedge \theta_3 + \tfrac{1}{16}\,(24\,\eta_1 + 40\,(\theta_{22} - \xi_2) - 9\,\xi_3) \wedge \theta_{13},
\end{aligned}
\tag{35}
$$

$$
\begin{aligned}
d\theta_{22} &= \eta_3 \wedge \xi_1 + \tfrac{1}{16}\,(8\,(\eta_2 - \theta_{22}) + 3\,\xi_3) \wedge \xi_2 - \eta_2 \wedge \xi_3,\\
d\theta_{23} &= \tfrac{1}{2}\,\eta_1 \wedge \theta_{23} + \eta_2 \wedge (\theta_{22} - \xi_2) + \eta_4 \wedge \xi_1 + \tfrac{1}{64}\,(8\,(\theta_{22} - \xi_2) + (64\,V - 1)\,\xi_3) \wedge \theta_2\\
&\quad + \tfrac{1}{8}\,(8\,(\eta_3 - \theta_3) - 5\,\theta_0 + 12\,U\,\xi_2) \wedge \xi_3 + \tfrac{1}{8}\,\theta_{22} \wedge (3\,(4\,\theta_{23} - \xi_2) - 8\,U\,\xi_3)\\
&\quad + \tfrac{1}{16}\,\theta_{23} \wedge (24\,\xi_2 - 7\,\xi_3)\,,\\
d\eta_1 &= -\tfrac{1}{8}\,(2\,\eta_2 + 6\,\theta_2 - 16\,(V\,\theta_{22} + \theta_{23}) + 4\,(2\,V - 1)\,\xi_2 + (8\,V - 3\,U)\,\xi_3) \wedge \xi_1\\
&\quad - \tfrac{5}{8}\,(\theta_{22} - \xi_2) \wedge \xi_3,\\
d\eta_2 &= \tfrac{1}{16}\,(8\,(\eta_1 + \theta_{22} - U\,\xi_1 - \xi_2) + 3\,\xi_3) \wedge \eta_2 + \tfrac{5}{16}\,\theta_0 \wedge \xi_1 - \tfrac{1}{8}\,\theta_2 \wedge (3\,V\,\xi_1 - \xi_3)\\
&\quad + \tfrac{1}{2}\,\theta_3 \wedge \xi_1 + (\theta_{12} + V\,\theta_{23} + V\,(V + U)\,\xi_3) \wedge \xi_1 + \tfrac{1}{2}\,(\theta_{22} + (U - 2\,V)\,\xi_1) \wedge \xi_2\\
&\quad + \big(2\,V\,\theta_{22} + \theta_{23} - \tfrac{1}{4}\,(8\,V + 1)\,\xi_2\big) \wedge \xi_3,\\
d\eta_3 &= \tfrac{1}{16}\,\eta_1 \wedge (8\,(\eta_3 + V\,\theta_2 + \theta_3) + 5\,\theta_0 - 2\,U\,\xi_2) - \theta_{12} \wedge \big(\big(V + \tfrac{9}{8}\big)\,\xi_1 + \xi_3\big) - \theta_{13} \wedge \xi_1\\
&\quad + \tfrac{1}{2}\,\eta_2 \wedge (2\,\theta_2 + 2\,(U - 2\,V)\,\xi_1 - \xi_2 + U\,\xi_3) + \eta_4 \wedge \xi_1 - \tfrac{5}{8}\,\theta_1 \wedge \xi_1\\
&\quad + \tfrac{1}{16}\,(24\,\theta_{22} + 8\,(3\,U - 8\,V)\,\xi_1 - 16\,\xi_2 - 5\,\xi_3) \wedge \eta_3 + \tfrac{1}{32}\,(19\,U - 8\,V)\,\xi_2 \wedge \xi_3\\
&\quad + \tfrac{5}{128}\,\theta_0 \wedge (8\,\theta_{22} + 16\,(U - 4\,V - 1)\,\xi_1 - 11\,\xi_3) - \tfrac{1}{2}\,\big(U^2 - 2\,U\,V - 8\,V^2\big)\,\xi_1 \wedge \xi_3\\
&\quad + \tfrac{1}{16}\,\theta_2 \wedge (8\,V\,\theta_{22} + 2\,((16\,V + 9)\,U - (56\,V + 9)\,V)\,\xi_1 - 9\,\xi_2 + 3\,V\,\xi_3)\\
&\quad + \tfrac{1}{16}\,\theta_3 \wedge (8\,\theta_{22} + 16\,(2\,U - 4\,V - 1)\,\xi_1 - 11\,\xi_3)\\
&\quad + \theta_{22} \wedge \big(V\,(U - 12\,V)\,\xi_1 + \big(\tfrac{1}{4}\,U + V\big)\,\xi_2\big)\\
&\quad + \tfrac{1}{2}\,\theta_{23} \wedge ((U - 2\,V)\,\xi_1 - \xi_2 - 2\,V\,\xi_3) + \tfrac{1}{2}\,\big((U - 2\,V)\,(U - 1) - 24\,V^2\big)\,\xi_1 \wedge \xi_2,\\
d\eta_4 &= \eta_8 \wedge \xi_1 + \tfrac{1}{32}\,(60\,\theta_0 + (96\,V - 1)\,\theta_2 + 96\,\theta_3 + 48\,U\,\theta_{22} - 24\,\theta_{23} - 96\,U\,\xi_2\\
&\quad +32\,(2\,V - U)\,\xi_3) \wedge \eta_2 + \tfrac{1}{8}\,(24\,\eta_2 + 7\,\theta_2 - 12\,\theta_{23} + 3\,\xi_2 + 4\,(7\,U - 8\,V)\,\xi_3) \wedge \eta_3\\
&\quad + \big(\eta_1 + 3\,(\theta_{22} - \xi_2) - \tfrac{7}{4}\,\xi_3\big) \wedge \eta_4 + \tfrac{1}{16}\,(17\,U\,\theta_{22} + (24\,V - 1)\,\theta_{23} - 6\,(V + 4\,U)\,\xi_2) \wedge \theta_2\\
&\quad - \tfrac{5}{64}\,(7\,(\theta_2 - \theta_{22}) + 12\,\theta_{23} + 4\,(\xi_2 - (7\,U - 8\,V - 2)\,\xi_3)) \wedge \theta_0\\
&\quad + \tfrac{1}{8}\,\theta_3 \wedge (7\,(\theta_2 + \theta_{22}) - 12\,\theta_{23} - 4\,\xi_2 - 8\,(7\,U - 8\,V - 2)\,\xi_3)\\
&\quad - \tfrac{1}{64}\,(72\,(\theta_{22} + \xi_2) + 73\,\xi_3) \wedge \theta_{12} + \tfrac{1}{128}\,\big(U\,(448\,V + 143) - 16\,V\,(32\,V^2 + 9)\big)\,\theta_2 \wedge \xi_3\\
&\quad + \big(\big(V - \tfrac{11}{16}\,U\big)\,\xi_2 - (U^2 - 13\,V^2)\,\xi_3\big) \wedge \theta_{22} + \tfrac{1}{8}\,(2\,(3\,U + 1)\,\xi_2 + (U + 4\,V)\,\xi_3) \wedge \theta_{23}\\
&\quad + \tfrac{1}{4}\,(4\,V\,(13\,V^2 - 1) - U\,(11\,U - 8\,V - 2))\,\xi_2 \wedge \xi_3,\\
d\eta_5 &= \tfrac{3}{2}\,\eta_1 \wedge \eta_5 + \eta_2 \wedge \big(4\,\eta_4 + \tfrac{1}{2}\,\theta_{12} - 2\,U\,\theta_{23} + (U - 2\,V)\,\xi_2\big) + \big(2\,\theta_2 - \xi_2 + \tfrac{5}{2}\,U\,\xi_3\big) \wedge \eta_4\\
&\quad + \tfrac{1}{16}\,(56\,(\theta_{22} - \xi_2) - 13\,\xi_3) \wedge \eta_5 + \eta_8 \wedge \xi_3 + \eta_9 \wedge \xi_1 + \tfrac{5}{8}\,\theta_1 \wedge (\theta_{22} - \xi_2)\\
&\quad + \tfrac{5}{64}\,(16\,\eta_3 - (24\,V + 1)\,\theta_2 + 16\,\theta_{12} - 8\,U\,(\theta_{22} + \theta_{23} + (U - 4\,V + 1)\,\xi_2)) \wedge \theta_0\\
&\quad + \tfrac{1}{16}\,\theta_2 \wedge (16\,V\,\eta_3 - 2\,(24\,V + 1)\,\theta_3 - 8\,(4\,V + 1)\,\theta_{12} - 48\,V\,(U - 2\,V)\,\theta_{22}\\
&\quad -(U + V)\,\theta_{23} + 2\,(2\,U\,(2\,V + 5) + V\,(16\,V - 11))\,\xi_2 + (U^2 - 4\,U\,V - 4\,V^2)\,\xi_3\big)\\
&\quad + (2\,(\eta_3 + \theta_{12}) - \theta_{23} + (4\,V - 2\,U + 1)\,\xi_2) \wedge \theta_3 + \theta_{13} \wedge (\theta_{22} - \xi_2)\\
&\quad + \tfrac{1}{8}\,(16\,(\eta_3 + 2\,V\,\theta_{22} - \theta_{23}) + (4\,U - 16\,V + 13)\,\xi_2 - 4\,(4\,V - 3\,U)\,\xi_3) \wedge \theta_{12}\\
&\quad + \big(\tfrac{1}{2}\,U^2 + U\,V - 12\,V^2\big)\,\theta_{22} \wedge \xi_2 + \tfrac{1}{2}\,\big(3\,(2\,V - U)\,\xi_2 + \big(4\,V\,(U + V) - 3\,U^2\big)\,\xi_3\big) \wedge \theta_{23}\\
&\quad - \tfrac{1}{2}\,\big(2\,(3\,U - 4\,V)\,\eta_3 + \big(8\,V^2 + 2\,U\,V - U^2\big)\,\xi_3\big) \wedge \xi_2,\\
d\eta_6 &= \tfrac{1}{64}\,\eta_2 \wedge (320\,\eta_5 + (80\,V - 85\,U)\,\theta_0 - 60\,\theta_1 + 128\,V\,(U + V)\,\theta_2 + 24\,U\,\theta_3 - 160\,U\,\theta_{12})
\end{aligned}
$$

$$
\begin{aligned}
&+ \big(6\,V\,\theta_3 - \tfrac{5}{2}\,\theta_{13}\big) \wedge \eta_3 + \eta_4 \wedge (V\,\theta_2 + 2\,(\theta_3 - U\,\xi_2)) - 3\,\eta_5 \wedge (\theta_2 + U\,\xi_3) \\
&+ \big(2\,\eta_1 + 4\,\theta_{22} - 4\,\xi_2 - \tfrac{7}{8}\,\xi_3\big) \wedge \eta_6 + \eta_8 \wedge \xi_2 + \eta_9 \wedge \xi_3 + \eta_{10} \wedge \xi_1 \\
&+ \tfrac{5}{64}\,\big(8\,(\eta_4 + (5\,U - 4\,V)\,\eta_3) + 10\,\theta_1 - 2\,\theta_{12} - 4\,(5\,U + 4\,V - 2)\,\theta_3 + 2\,(7\,V - 8\,U)\,\theta_{23} \\
&-(96\,V^2 + 7\,U - 10\,U^2 + 8\,U\,V - 14\,V)\,\wedge \xi_2 + 2\,(17\,V^2 - 5\,U^2 + 7\,U\,V)\,\xi_3\big) \wedge \theta_0 \\
&+ \tfrac{5}{128}\,(32\,(\eta_3 - \theta_3) + 8\,(U\,\theta_{22} - \theta_{23} + U\,\xi_2) - 9\,U\,\xi_3) \wedge \theta_1 \\
&+ \tfrac{1}{256}\,((U\,(400\,V + 310) - V\,(640\,V + 295))\,\theta_0 + 40\,(8\,V + 5)\,\theta_1) \wedge \theta_2 \\
&+ \tfrac{1}{32}\,\big((14\,U + 192\,V^2 + 23\,V)\,\theta_2 - 52\,\theta_{12} + 288\,V\,(2\,V - U)\,\theta_{22} + 8\,(3\,V - 2\,U)\,\theta_{23} \\
&+4\,(3\,U\,(16\,V - 1) - 6\,V\,(24\,V + 1))\,\xi_2 + 24\,V\,(V + U)\,\xi_3) \wedge \theta_3 \\
&- \tfrac{5}{64}\,(8\,(\theta_{22} + \xi_2) - 3\,\xi_3) \wedge \theta_{11} + \theta_{12} \wedge \big(\big(U + \tfrac{9}{8}\,V\big)\,\theta_2 - 3\,V\,\theta_{23} + 2\,(U - 2\,V)\,\xi_2 \\
&+2\,(U^2 - 2\,V^2 - 2\,U\,V)\,\xi_3\big) + \tfrac{1}{16}\,\theta_{13} \wedge (25\,\theta_0 + 40\,(\theta_3 + \theta_{23}) + (40\,V + 23)\,\theta_2 \\
&-144\,V\,\theta_{22} + 4\,(36\,V - 5\,U - 3)\,\xi_2 + 12\,(2\,V - U)\,\xi_3) + \tfrac{5}{8}\,V\,(U - 12\,V)\,\theta_0 \wedge \theta_{22} \\
&+ \tfrac{3}{4}\,U\,(2\,V\,\theta_2 - U\,\xi_2) \wedge \theta_{23} + \tfrac{1}{2}\,V\,\theta_2 \wedge ((U - 6\,V)\,\xi_2 - V\,(5\,U + 12\,V)\,\xi_3)\,,
\end{aligned}
$$

$$
\begin{aligned}
d\eta_7 \ =\ & 3\,\eta_2 \wedge \big(2\,\eta_6 - \tfrac{1}{4}\,\theta_{11} - \theta_{13}\big) - \eta_3 \wedge (13\,V\,\theta_1 - 3\,\theta_{11}) + 2\,\eta_4 \wedge (\theta_1 - V\,\theta_3) \\
&+ \tfrac{1}{8}\,\eta_5 \wedge (19\,\theta_0 + 16\,(V\,\theta_2 + \theta_3) - 20\,U\,\xi_2) - \tfrac{1}{2}\,\eta_6 \wedge (8\,\theta_2 + 7\,U\,\xi_3) \\
&+ \tfrac{1}{16}\,(40\,\eta_1 + 72\,(\theta_{22} - \xi_2) - 15\,\xi_3) \wedge \eta_7 - \eta_8 \wedge \theta_0 + \eta_9 \wedge \xi_2 + \eta_{10} \wedge \xi_3 + \eta_{11} \wedge \xi_1 \\
&+ \tfrac{1}{32}\,\big(4\,(13\,U^2 - 248\,V^2 + 4\,U\,V)\,\eta_2 - 256\,V\,(U + V)\,\eta_3 + 8\,(6\,U + 7\,V)\,\eta_4 \\
&-5\,(52\,V + 1)\,\theta_1 + (128\,V^3 + V^2\,(4\,U - 345) + 56\,U^2 + 45\,U\,V)\,\theta_2 \\
&+8\,(16\,V\,(U + V) - 3\,U)\,\theta_3 + 60\,\theta_{11} - (88\,U + 59\,V)\,\theta_{12} - 72\,\theta_{13} \\
&-32\,V\,(U^2 - 2\,U\,V - 12\,V^2)\,\theta_{22} + 12\,V\,(U + 2\,V)\,\theta_{23} \\
&-2\,(192\,V^3 + U^2\,(V - 11) + 8\,U\,V + 4\,(U - 3)\,V^2)\,\xi_2 \\
&-2\,(92\,U\,V^2 - 15\,U^3 + 6\,U^2\,V + 360\,V^3)\,\xi_3\big) \wedge \theta_0 + U^2\,\theta_{12} \wedge \xi_2 \\
&+ \tfrac{1}{16}\,(16\,(3\,U - 4\,V)\,\eta_2 + 4\,(52\,V - 3)\,\theta_3 - 36\,\theta_{12} - 16\,V\,(20\,U - 39\,V)\,\theta_{22} \\
&+28\,V\,\theta_{23} + 2\,(3\,U\,(36\,V - 1) - 2\,V\,(156\,V - 7))\,\xi_2 \\
&+ \big(17\,(U^2 - 4\,V^2) + 12\,U\,V\big)\ \xi_3\big) \wedge \theta_1 \\
&+ \tfrac{1}{32}\,\big(\big(117\,V - 416\,V^2 - 88\,U\big)\,\theta_1 + 4\,(24\,V + 23)\,\theta_{11} - 32\,V\,(U + 3\,V)\,\theta_{12} \\
&+8\,(4\,U + 9\,V)\,\theta_{13}) \wedge \theta_2 + \tfrac{1}{4}\,(16\,V\,(U + V)\,\eta_2 + 9\,V\,(2\,U - V)\,\theta_2 + 12\,\theta_{11} \\
&-(4\,U + 3\,V)\,\theta_{12} - 4\,\theta_{13} - 12\,V\,(U - 2\,V)\,\theta_{23} + 2\,(U^2 + 4\,U\,V + 12\,V^2)\,\xi_2 \\
&-4\,V\,(U^2 - 4\,U\,V - 12\,V^2)\,\xi_3\big) \wedge \theta_3 + \tfrac{3}{8}\,(40\,V\,\theta_{22} - 8\,\theta_{23} + 4\,(U - 10\,V + 1)\,\xi_2 \\
&+(9\,U - 8\,V)\,\xi_3) \wedge \theta_{11} + (6\,V\,\theta_{23} - 3\,(U - 2\,V)\,\xi_2 \\
&+ \big(6\,V\,(U + V) - \tfrac{5}{2}\,U^2\big)\ \xi_3\big) \wedge \theta_{13}.
\end{aligned}
$$

Bibliography

[1] O.I. Bogoyavlenskiy, Breaking Solitons. Nonlinear Integrable Equations. Moscow, Nauka (1991)

[2] Bryant, R.L., Chern, S.S., Gardner, R.B., Goldschmidt, H.L., Griffiths, P.A.: Exterior Differential Systems. N.Y., Springer-Verlag (1991)

[3] Bryant, R.L., Griffiths, Ph.A.: Characteristic cohomology of differential systems (II): conservation laws for a class of parabolic equations, Duke Math. J. **78**, 531–676 (1995)

[4] Cartan, É.: Sur la structure des groupes infinis de transformations. Œuvres Complètes, Part II, **2**, p. 571–715. Gauthier - Villars, Paris (1953)

[5] Cartan, É.: Les sous-groupes des groupes continus de transformations. Œuvres Complètes, Part II, **2**, p. 719–856. Gauthier - Villars, Paris (1953)

[6] Cartan, É.: Les problèmes d'équivalence. Œuvres Complètes, Part II, **2**, p. 1311–1334. Gauthier - Villars, Paris (1953)

[7] Cartan, É.: La structure des groupes infinis. Œuvres Complètes, Part II, **2**, p. 1335–1384. Gauthier - Villars, Paris (1953)

[8] Dunajski, M.: A class of Einstein–Weil spaces associated to an integrable system of hydrodynamic type, J. Geom. Phys. **51**, 126-137 (2004)

[9] Dunajski, M.: An interpolating dispersionless integrable system. J. Phys. A, Math. Theor., **41**, 315202 (2008)

[10] Gardner, R.B.: The method of equivalence and its applications. CBMS–NSF regional conference series in applied math., SIAM, Philadelphia (1989)

[11] Fels, M., Olver, P.J.: Moving coframes. I. A practical algorithm. Acta. Appl. Math. **51**, 161–213 (1998)

[12] Kamran, N.: Contributions to the Study of the Equivalence Problem of Élie Cartan and its Applications to Partial and Ordinary Differential Equations. Mem. Cl. Sci. Acad. Roy. Belg., **45**, Fac. 7 (1989)

[13] Zabolotskaya, E.A., Khokhlov, R.V.: Quasi-plane waves in the nonlinear acoustics of confined beams. Sov. Phys. Acoust., **15**, 35–40 (1969)

[14] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal symmetries and the theory of coverings. Acta Appl. Math., **2**, 79–86 (1984)

[15] Krasil'shchik, I.S., Lychagin, V.V., Vinogradov, A.M.: Geometry of Jet Spaces and Nonlinear Partial Differential Equations. Gordon and Breach, New York (1986)

[16] Krasil'shchik, I.S., Vinogradov, A.M.: Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. Acta Appl. Math., **15**, 161–209 (1989)

[17] Krasil'shchik, I.S., Vinogradov, A.M. (eds.): Symmetries and Conservation Laws for Differential Equations of Mathematical Physics. Transl. Math. Monographs 182, Amer. Math. Soc., Providence (1999).

[18] Marvan, M.: On zero-curvature representations of partial differential equations. Proc. Conf. on Diff. Geom. and Its Appl., Opava (Czech Republic), 103–122 (1992)

[19] Morozov, O.I.: Moving coframes and symmetries of differential equations. J. Phys. A, Math. Gen., **35**, 2965–2977 (2002)

[20] Morozov, O.I.: Contact-equivalence problem for linear hyperbolic equations. J. Math. Sci., **135**, 2680–2694 (2006)

[21] Morozov, O.I.: Cartan's structure theory of symmetry pseudo-groups, coverings and multi-valued solutions for the Khokhlov–Zabolotskaya equation, Acta Appl. Math. **101**, 231-241 (2008)

[22] Morozov, O.I. Contact integrable extensions of symmetry pseudo-group and coverings of the r-th modified dispersionless Kadomtsev–Petviashvili equation. `arXiv:0809.1218v1 [math.DG]` (2008)

[23] Olver, P.J.: Equivalence, Invariants, and Symmetry. Cambridge, Cambridge University Press (1995)

[24] Pavlov, M.V. Integrable hydrodynamic chains. J. Math. Phys., **44**, 4134–4156 (2003)

[25] Plebañski, J.F. Some solutions of complex Einstein equations. J. Math. Phys., **16**, 2395–2402 (1975)

[26] Stormark, O.: Lie's Structural Approach to PDE Systems. Cambridge, Cambridge University Press (2000)

[27] Vasil'eva, M.V.: The Structure of Infinite Lie Groups of Transformations. Moscow, MGPI (1972) (in Russian)

[28] Wahlquist, H.D., Estabrook F.B.: Prolongation structures of nonlinear evolution equations. J. Math. Phys., **16**, 1–7 (1975)